




Changing Northern Landscapes



Maps of Selected NASA ABoVE Research
Bristol Bay, Alaska



Meet the Interns

Behind the Pages:

During her internship, Beverly Bolster was a senior at the University of Maryland, College Park. She pursued a dual degree in Geospatial Data Science and Sociology, with minors in Spanish and Sustainability Studies. She was passionate about community engagement to promote sustainability and address the impacts of climate change, and she started this booklet during her internship with NASA Arctic-Boreal Vulnerability Experiment. She planned to work at a summer program for children with severe emotional and behavior challenges before pursuing additional experience working with children, families, and communities and ultimately attending graduate school for developmental psychology or social work.



Mary Banner is Ponca and an enrolled member of the Mille Lacs Band of Ojibwe. She is kahu to four rescued pups and celebrated ten years sober. She graduated with a Master of Professional Studies in Environmental Science from State University of New York College of Environmental Science and Forestry with a focus on ecosystems: land, air, and water. This booklet completed her internship with NASA Arctic-Boreal Vulnerability Experiment, and her hope is that these one-pagers showing complex data are more accessible to communities. She enjoys spending time outdoors, exploring with her animals, watching sports, and seeking opportunities to work with Tribal Nations, state, local, and within the private sector. Hy'shqe (Thank you).



Table of Contents

Introduction	1
Land Cover Change	2
<i>ABoVE: Landsat-derived Annual Dominant Land Cover Across ABoVE Core Domain, 1984-2014</i>	
Bristol Bay	3
Areas Burned by Wildfire	4
<i>Arctic Boreal Annual Burned Area, Circumpolar Boreal Forest and Tundra, V2 2002-2022</i>	
Bristol Bay	5
Ground Thaw Depth	6
<i>ABoVE: Active Layer Thickness from Airborne L- and P- band SAR, Alaska, Ver. 3</i>	
Bristol Bay	7
Wildfire Burn Area and Severity	8
<i>ABoVE: Burned Area, Depth, and Combustion for Alaska and Canada, 2001-2019</i>	
Bristol Bay	9
Forest Biomass (Carbon Storage)	10
<i>Aboveground Biomass Density for High Latitude Forests from IceSAT2</i>	
Bristol Bay	11
Wildfires	12
<i>ABoVE: Ignitions of Above-Fire Emission Database Fires in Alaska and Canada</i>	
Bristol Bay	13
Projected Changes in Land Cover	14
<i>Land Use and Land Cover Change Projection in the ABoVE Domain</i>	
Bristol Bay	15
Forest Canopy Cover and Forest Age	16
<i>ABoVE: Tree Canopy Cover and Stand Age from Landsat, Boreal Forest Biome, 1984-2020</i>	
Bristol Bay	17
Carbon Dioxide Absorption and Release	18
<i>Machine learning-based Arctic-boreal terrestrial ecosystem CO₂ Fluxes, 2001-2020</i>	
Bristol Bay	19
Carbon Dioxide and Methane	20
<i>Gridded CO₂ and CH₄ Flux Estimates for pan-Arctic and Boreal Regions, 2003-2015</i>	
Bristol Bay	21
Conclusion	22
Resources, Credits, & Acknowledgments	23

Introduction

This booklet presents a variety of datasets created through NASA's Arctic-Boreal Vulnerability Experiment (ABoVE). ABoVE is a NASA Terrestrial Ecology Program field campaign being conducted in Alaska and western Canada which began in 2015. It is a diverse large-scale study of the impacts of environmental change on arctic and boreal terrestrial and freshwater ecosystems, and the implications of these changes for social and ecological systems. It encompasses the variability in the key types of ecosystems that are both unique to Arctic and boreal regions in North America as well as being representative of the larger Northern High Latitude region.

We aim to make the data more widely available and accessible to communities, governments, and other organizations.

For each type of data, there is a page with a general explanation and a map of the full extent that the data covers. We also provide the following information:

What do the data show?

This section explains the importance of the dataset, as well as the type of information that can be interpreted from the maps provided.

How were the data produced?

This section explains the methods for capturing the data. This may include whether the data were collected through airborne campaigns (in which sensors attached to aircraft gather information about the earth's surface) or satellites (which contain instruments to collect images as they orbit the earth). Additionally, this may explain models or algorithms that were used to produce data.

Spatial Resolution:

This includes a value that tells the distance between each measurement that is recorded. Smaller distances lead to a higher level of detail, but less area can be covered in the same amount of time.

Spatial Coverage:

This details the regional extent that the dataset includes.

Temporal Resolution:

This tells how often the data were measured, whether this occurred only once or how frequently measurements recurred.

Temporal Coverage:

This includes a data range that tells the time period that the dataset covers.

Citation


There is also a citation with a link to further information and a platform to download each dataset.

Regional Example Maps

Where applicable, additional maps for regions in Alaska and Canada are provided. These show how the data can be visualized at a community or local level.

Land Cover Change

Dataset: ABoVE: Landsat-derived Annual Dominant Land Cover Across ABoVE Core Domain, 1984-2014



Land cover across ABoVE core domain 2014

What do the data show?

These data include land cover classifications over Alaska and western Canada from 1984-2014. One dataset classifies land cover into fifteen classes of forest and shrub types, and the other simplifies this classification into ten categories.

How were the data produced?

The datasets were produced by determining the dominant plant type in each 30 meter by 30 meter pixel (square). Thus, each square is classified according to which plant type covered the largest area. The resulting maps provide a consistent view of vegetation patterns across Alaska and western Canada.

Citation:


Wang, J.A., D. Sulla-Menashe, C.E. Woodcock, O. Sonnentag, R.F. Keeling, and M.A. Friedl. 2019. ABoVE: Landsat-derived Annual Dominant Land Cover Across ABoVE Core Domain, 1984-2014. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/1691>

Spatial Resolution: 30 m

Spatial Coverage: Alaska and Canada


Temporal Resolution: Annual

Temporal Coverage: 01-01-1984 to 12-31-2014



Conifer trees represent the evergreen forest land cover, shown in dark green on the map. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)

Bristol Bay




Land cover in Bristol Bay Region 2014

Disclaimer: This dataset doesn't extend all the way across the peninsula

Areas Burned by Wildfire

Dataset: Arctic Boreal Annual Burned Area, Circumpolar Boreal Forest and Tundra, V2, 2002-2022



Burned area between 2002 and 2022

What do the data show?

The dataset provides the amount of area that was burned in boreal forests and tundra in the northern latitudes areas above 50 degrees north latitude each year from 2002 to 2022.

How were the data produced?

The data were produced using satellite data at 500m spatial resolution (MODIS data) through an algorithm that captures the difference before and after a fire to assess the amount of area burned. In particular, this algorithm was intended to capture fires that occur late in the season (near the end of the northern hemisphere summer months) and unburned areas within fire perimeters.

Citation:

Loboda, T.V., J.V. Hall, D. Chen, A. Hoffman-Hall, V.S. Shevade, F. Argueta, and X. Liang. 2024. Arctic Boreal Annual Burned Area, Circumpolar Boreal Forest and Tundra, V2, 2002-2022. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/2328>

Spatial Resolution: 463 m

Spatial Coverage: High northern latitudes (circumpolar above 50 degrees N)


Temporal Resolution: Annual

Temporal Coverage: 2002 to 2022



Burned trees following a wildfire. Areas shown in red on the map indicate locations that burned between 2002 and 2022. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)


Bristol Bay



Bristol Bay burned area between 2002 and 2022

Ground Thaw Depth

Dataset: ABoVE: Active Layer Thickness from Airborne L- and P- band SAR, Alaska, 2017, Ver. 3



Active layer thickness measurements collected along ABoVE flight paths in 2017

What do the data show?

This dataset estimates changes in the presence of permafrost and soil moisture by calculating active layer thickness, seasonal subsidence, and the vertical soil moisture profile at the time of predicted maximum thaw in 2017. The active layer thickness is the depth of the ground that thaws each summer. The seasonal subsidence is how much the Earth's surface moves downward each year due to the ground becoming drier. The vertical soil moisture profile is how much the quantity of water in the soil changes as the depth of the soil changes, which impacts how quickly water can pass through soil and how quickly soil will dry out.

How were the data produced?

These data were produced through NASA's ABoVE airborne campaign. Sensors attached to aircraft gathered images of the earth's surface above Alaska and Canada. The data product was created by the Permafrost Dynamics Observatory (PDO) project.

Citation:

Chen, R.H., R.J. Michaelides, J. Chen, A.C. Chen, L.K. Clayton, K. Bakian-Dogaheh, L. Huang, E. Jafarov, L. Liu, M. Moghaddam, A.D. Parsekian, T.D. Sullivan, A. Tabatabaenejad, E. Wig, H.A. Zebker, and Y. Zhao. 2022. ABoVE: Active Layer Thickness from Airborne L- and P- band SAR, Alaska, 2017, Ver. 3. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORN LDAAC/2004>

Spatial Resolution: 30 m

Spatial Coverage: 51 sites across the ABoVE domain, including 39 in Alaska and 12 in Canada; Alaska, Yukon, Northwest Territories Yukon, Northwest Territories


Temporal Resolution: One time estimate in 2017

Temporal Coverage: 06-19-2017 to 09-16-2017



Soil from permafrost terrain. Active layer thickness measures how deeply the ground thaws each summer. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE).


Bristol Bay



Bristol Bay active layer thickness 2017

Wildfire Burn Area and Severity

Dataset: ABoVE: Burned Area, Depth, and Combustion for Alaska and Canada, 2001-2019



Total carbon emissions combustion between 2001-2019 aggregated to a 70 km grid

What do the data show?

This dataset shows fire activity in Alaska and Canada from 2001 to 2019, showing where and when fires occurred, how much area burned, and the amount of carbon released into the atmosphere. It uses satellite data, field observations, and advanced modeling to help understand the impacts of wildfires on boreal ecosystems, including carbon emissions and ecosystem disturbances.

How were the data produced?

The data were produced using MODIS and Landsat satellite imagery to detect fire locations and burned areas, combined with statistical modeling of carbon combustion and burn depth based on field observations and environmental variables like fire severity, topography, and climate.

Citation:

Potter, S., S. Veraverbeke, X.J. Walker, M.C. Mack, S.J. Goetz, J.L. Baltzer, C. Dieleman, N.H.F. French, E.S. Kane, M.R. Turetsky, E.B. Wiggins, and B.M. Rogers. 2022. ABoVE: Burned Area, Depth, and Combustion for Alaska and Canada, 2001-2019. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAC/2063>

Spatial Resolution: 500 m

Spatial Coverage: Alaska and Canada


Temporal Resolution: Annual

Temporal Coverage: 01-01-2001 to 12-31-2019



Wildfire burn severity can vary across the landscape, leaving a mix of heavily and lightly burned areas. Photo courtesy of Dr. Elizabeth Hoy, NASA ABoVE


Bristol Bay



Bristol Bay total fire combustion 2001 to 2019 measured in kg C

Carbon Exchange Between Land and Atmosphere

Dataset: Aboveground Biomass Density for High Latitude Forests from IceSAT2



What do the data show?

This dataset shows how much woody biomass is stored in Northern forests, helping to map and monitor forest carbon stocks.

How were the data produced?

The data were created by combining ground measurements with satellite laser data and advanced models to estimate biomass across these forests from 2019 to 2021.

Citation:


Duncanson, L., P.M. Montesano, A. Neuenschwander, N. Thomas, A. Mandel, D. Minor, E. Guenther, S. Hancock, T. Feng, A. Barciauskas, G.W. Chang, S. Shah, and B.P. Satorius. 2023. Aboveground Biomass Density for High Latitude Forests from ICESat-2, 2020. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/2186>

Spatial Resolution: 30 m

Spatial Coverage: Boreal forest zone in the Northern Hemisphere, covering areas above 44 degrees latitude where these forests thrive

Temporal Resolution: Circa 2020 (06-01-2019 to 09-30-2021)

Temporal Coverage: One-time measurements




Bristol Bay



Aboveground biomass (Mg ha^{-1}) for boreal forest estimated from ICESat-2 imagery

Wildfires

Dataset: ABoVE: Ignitions of Above-Fire Emission Database Fires in Alaska and Canada



Fire Ignition locations across Canada and Alaska, U.S., for 2001 - 2019

What do the data show?

The dataset shows where and when wildfires started in boreal forests across Alaska and Canada from 2001 to 2019.

How were the data produced?

The data was produced by analyzing satellite images to detect fire activity, using specialized methods to identify ignition points and determine their timing within a single day.

Citation:

Hessilt, T.D., B.M. Rogers, R.C. Scholten, S. Potter, T.A.J. Janssen, and S. Veraverbeke. 2023. ABoVE: Ignitions of ABoVE-FED Fires in Alaska and Canada. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/2316>

Spatial Resolution: Points with 462 Location precision

Spatial Coverage: Alaska and Canada


Temporal Resolution: Daily

Temporal Coverage: 01-01-2001 to 12-31-2019



A boreal forest following wildfire, photographed during NASA ABoVE field research. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)



Bristol Bay



Bristol Bay Fire Ignitions, 2001 - 2019

Projected Changes in Land Cover

Dataset: Land Use and Land Cover Change Projection in the ABoVE Domain



Percentage Cover of Broadleaf Deciduous Boreal Tree predicted under scenario SSP 1 - 2.6 in 2015 and 2100

What do the data show?

The data shows predicted changes in land use and land cover in the Arctic region from 2015 to 2100, including shifts in forests, shrubs, grasses, and crops. The maps above specifically highlight percent cover of deciduous boreal trees under climate scenario SSP 1 - 2.6 in 2015 and 2100.

How were the data produced?

The data was produced by combining global land use model with satellite data and to create detailed annual maps under two different climate and socioeconomic scenarios which model differences in human-driven land use changes and diverse land cover types.

Citation:

Luo, M., F. Li, D. Hao, Q. Zhu, H. Dashti, and M. Chen. 2024. Land Use and Land Cover Change Projection in the ABoVE Domain. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/2353>

Spatial Resolution: 0.25 degrees

Spatial Coverage: ABoVE domain, Alaska and Canada

Temporal Resolution: Annual



Temporal Coverage: 01-01-2015 to 12-31-2100



Broadleaf deciduous trees, highlighted in the maps above, may become more widespread across parts of the Arctic-Boreal region under future climate scenarios. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)


Bristol Bay

Predicted Land Cover in Broadleaf Deciduous Boreal Tree under climate scenario SSP 1 - 2.6 in 2015 and 2100.



Forest Canopy Cover and Forest Age

Dataset: ABoVE: Tree Canopy Cover and Stand Age from Landsat, Boreal Forest Biome, 1984-2020



Tree Canopy Cover in the ABoVE Domain, 2020

What do the data show?

This dataset shows the amount of the circumpolar boreal forest covered by tree canopy and the age of the forests. Using this information, forest change estimates of stand age were made using data from 1984 to 2020.

How were the data produced?

The data were produced using satellite images from Landsat and machine learning to estimate tree canopy cover, then changes were tracked over time to determine changes in forest age.

Citation:

Feng, M., Sexton, J. O., Wang, P., Channan, S., Montesano, P. M., Wagner, W., Wooten, M. R., & Neigh, C. S. (2022). ABoVE: Tree Canopy Cover and Stand Age from Landsat, Boreal Forest Biome, 1984-2020. ORNL Distributed Active Archive Center. <https://doi.org/10.3334/ORNLDAAAC/2012>

Spatial Resolution: Pan-boreal including all boreal forest or taiga ecoregions, as well as temperate conifer forests and tundra regions predominantly surrounded by boreal forest or taiga

Spatial Coverage: 30 m


Temporal Resolution: Annual

Temporal Coverage: 01-01-1984 to 12-31-2020



Looking upward through a boreal forest canopy. Tree canopy cover measures how much of the ground is covered by the crowns of trees. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)

Bristol Bay




Bristol Bay Tree Canopy Cover in 2020

Disclaimer: This dataset doesn't extend all the way across the peninsula

Carbon Dioxide Absorption and Release

Dataset: Machine learning-based Arctic-boreal terrestrial ecosystem CO₂ Fluxes, 2001-2020



ABoVE Domain CO₂ Fluxes showing the Net Ecosystem Exchange and Net Ecosystem Fire from 2001-2020

What do the data show?

The dataset show how Arctic and boreal ecosystems acted as carbon sinks—absorbing more CO₂ than they released—or as carbon sources—releasing more CO₂ than they absorbed—between 2001 and 2020.

How were the data produced?

The dataset was created by combining ground-based CO₂ measurements with satellite, weather, soil, and landscape data, into machine learning models to estimate carbon flux across the region.

Citation:

Virkkala, A-M., B.M. Rogers, J.D. Watts, K. Arndt, S. Potter, I. Wargowsky, and S. Natali. 2024. Machine learning-based Arctic-boreal terrestrial ecosystem CO₂ fluxes, 2001-2020. ORNL DAAC, Oak Ridge, Tennessee, USA.
<https://doi.org/10.3334/ORNLDAAC/2377>

Spatial Resolution: 1 km

Spatial Coverage: Circumpolar arctic and boreal regions (tundra and boreal biomes), >49 degrees north latitude


Temporal Resolution: Monthly

Temporal Coverage: 01-01-2001 to 12-31-2020



Wetlands can store large amounts of carbon while also exchanging carbon dioxide with the atmosphere. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)


Bristol Bay



Bristol Bay CO₂ Fluxes showing the Net Ecosystem Exchange and Net Ecosystem Fire from 2001-2020

Carbon Dioxide and Methane

Dataset: Gridded CO₂ and CH₄ Flux Estimates for Pan-Arctic and Boreal Regions, 2003-2015



Tracking Methane Emissions Across Arctic Wetlands in the ABoVE Domain showing July in 2003 and 2015

What do the data show?

The data comes from a computer model that estimates plant growth, soil breakdown, and greenhouse gas emissions. It uses satellite images and environmental data collected from over 60 monitoring sites.

How were the data produced?

The data was created using a computer model that estimates plant growth, how soil breaks down, and greenhouse gas emissions. This model is based on satellite images and environmental data from the monitoring sites.

Citation:

Farina, M.K., and J.D. Watts. 2022. Gridded CO₂ and CH₄ Flux Estimates for pan-Arctic and Boreal Regions, 2003-2015. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/2121>

Spatial Resolution: Pan-Arctic and Boreal Zone (>49 Deg N)

Spatial Coverage: 1 km


Temporal Resolution: Daily

Temporal Coverage: 01-01-2003 to 12-31-2015



Wetlands are one of the largest natural sources of methane in Arctic and boreal ecosystems. Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE)

Bristol Bay



Tracking Methane Emissions Across Arctic Wetlands in Bristol Bay showing July in 2003 and 2015

Conclusion

Over the past decade, the Arctic-Boreal Vulnerability Experiment (ABoVE) has brought together scientists, data analysts, government and community leaders, and Traditional and Indigenous knowledge holders to better understand the profound environmental changes unfolding across the Arctic and boreal regions. These collaborations have generated invaluable datasets spanning multiple science themes, including:

Carbon dynamics and fluxes, fire regimes and post-fire recovery, vegetation structure and aboveground biomass, permafrost thaw and soil carbon dynamics, subsistence resources and environmental conditions, hydrology and watershed dynamics, remote sensing and spectroscopy, land use and cover change, ecosystem disturbance and recovery, cryosphere dynamics and ice monitoring, human-environment interactions and climate impacts, ecosystem health and biodiversity.

These themes not only advance scientific understanding but also support planning and decision-making by local and Indigenous communities whose lands and livelihoods are directly impacted by these changes, as well as state/territorial and federal land managers with regional and national priorities.

Each region within the ABoVE domain tells a unique story - a story of changing landscapes, shifting ecosystems, and community resilience. Through the creation of regional one-pagers, we have distilled some of ABoVE's complex scientific findings into accessible formats that can help residents make informed decisions. These region-specific summaries provide a bridge between cutting-edge science and the lived experiences of local and Indigenous peoples, ensuring that knowledge generated over the last decade serves those most affected by environmental change.

As we look to the future, the legacy of ABoVE extends beyond datasets and maps created. It lives in the relationships built, the knowledge shared, and the commitment to ensuring that data generated in the Arctic and boreal regions serves the communities who call these lands home. Moving forward, we remain committed to upholding principles of Indigenous data sovereignty, fostering open-access science, and cultivating deeper partnerships that respect and honor the knowledge and perspectives of governments, rightsholders, and residents of the Arctic and boreal regions.

This work would not have been possible without the dedicated efforts of the ABoVE science teams, data analysts, and field researchers who have contributed their expertise to this endeavor. We also extend our deepest gratitude to the Indigenous communities across the Arctic and boreal regions who have generously shared their knowledge, experiences, and perspectives to ensure that these datasets reflect the realities of the land. Special thanks to the Sahtu Region, Vuntut Gwitchin Traditional Territory, Bristol Bay, Fairbanks, Dehcho, Yukon-Kuskokwim Delta, Whitehorse, Mackenzie Delta, and all other communities who have guided and enriched this work.

Resources, Credits, & Acknowledgments

Data Sources Featured in this Booklet

The maps and figures in this booklet are based on NASA Arctic-Boreal Vulnerability Experiment (ABoVE) datasets archived and distributed through the Oak Ridge National Laboratory Distributed Active Archive Center (ORNL DAAC).

ABoVE: Landsat-derived Annual Dominant Land Cover Across ABoVE Core Domain, 1984-2014
Arctic Boreal Annual Burned Area, Circumpolar Boreal Forest and Tundra, Version 2, 2002-2022
ABoVE: Active Layer Thickness from Airborne L- and P-band SAR, Alaska, Version 3
ABoVE: Burned Area, Depth, and Combustion for Alaska and Canada, 2001-2019
Aboveground Biomass Density for High Latitude Forests from ICESat-2
Land Use and Land Cover Change Projection in the ABoVE Domain
ABoVE: Tree Canopy Cover and Stand Age from Landsat, Boreal Forest Biome, 1984-2020
Machine Learning-based Arctic-boreal Terrestrial Ecosystem CO₂ Fluxes, 2001-2020
Gridded CO₂ and CH₄ Flux Estimates for Pan-Arctic and Boreal Regions, 2003-2015

Cover Images Credit

Top Photo: *Arctic-Boreal Landscape (Denali National Park, Alaska). Photo courtesy of Dr. Elizabeth Hoy, NASA Arctic-Boreal Vulnerability Experiment (ABoVE).*

Bottom Photo: *NASA Satellite View of Bristol Bay, Alaska. Photo courtesy of Mary Banner, NASA Worldview image using MODIS data from the Terra satellite, August 16, 2019.*

Explore ABoVE Information and Products

ABoVE Project Website: <https://above.nasa.gov>

NASA Earthdata ABoVE Search: <https://search.earthdata.nasa.gov/search?portal=above>

NASA Earthdata Search (all NASA Earth data products): <https://search.earthdata.nasa.gov>

NASA Oak Ridge National Laboratory Distributed Active Archive Center for Biogeochemical Dynamics (ORNL DAAC): <https://www.earthdata.nasa.gov/data/projects/above>